



Subspaces


Linear Algebra

Department of Computer Engineering

Sharif University of Technology

Hamid R. Rabiee rabiee@sharif.edu

Maryam Ramezani maryam.ramezani@sharif.edu



Subspace Definition




Definition

A **non-empty subset** of vector space for which closure holds for addition and scalar multiplication is called a subspace.

Subspace: If V is a vector space and **subset** $U \subseteq V$, then **U is itself a vector space** with the **same** addition and scalar multiplication as V .

- Zero vector is a subspace of every vector space.
- Vector space is a subspace of itself.






A subspace of \mathbb{R}^n is any set H in \mathbb{R}^n that has these properties:

- The zero vector is in H .
- For each u and v in H , the sum $u + v$ is in H .
- For each u in H and each scalar c , the vector cu is in H .

Example

- $H = \text{Span} \{x_1, x_2\}$, then H is a subspace of \mathbb{R}^2 .

- Is L subspace of \mathbb{R}^2 ?



- The vector space \mathbb{R}^2 is a subspace of \mathbb{R}^3 ?

- Is H a subspace of \mathbb{R}^3 ? $H = \left\{ \begin{bmatrix} s \\ t \\ 0 \end{bmatrix} : s \text{ and } t \text{ are real} \right\}$

Vector Space vs Subspace



Let V be a vector space and let $U \subseteq V$:

Vector Space


1. $u + v \in V$
2. $u + v = v + u$
3. $(u + v) + w = u + (v + w)$
4. There is a vector $0 \in V$ such that $u + 0 = u$
5. For each $u \in V$, there is a vector $-u \in V$ such that $u + (-u) = 0$
6. $cu \in V$
7. $c(u + v) = cu + cv$
8. $(c + d)u = cu + du$
9. $c(du) = (cd)u$
10. $1u = u$

A subspace of \mathbb{R}^n is any set H in \mathbb{R}^n that has these properties:

- The zero vector is in H .
- For each u and v in H , the sum $u + v$ is in H .
- For each u in H and each scalar c , the vector cu is in H .

Subspace

1. $u + v \in U$
2. $u + v = v + u$
3. $(u + v) + w = u + (v + w)$
4. There is a vector $0 \in U$ such that $u + 0 = u$
5. For each $u \in U$, there is a vector $-u \in U$ such that $u + (-u) = 0$
6. $cu \in U$
7. $c(u + v) = cu + cv$
8. $(c + d)u = cu + du$
9. $c(du) = (cd)u$
10. $1u = u$







Theorem

A non-empty subset U of V is a subspace of V if and only if for each pair of vectors b, c in U and each scalar α in F the vector $\alpha b + c$ is again in U .

Proof:

Example

- In F^n , the set of n -tuples $\begin{bmatrix} x_1 \\ \dots \\ x_n \end{bmatrix}$ with $x_1 = 0$
- In F^n , the set of n -tuples $\begin{bmatrix} x_1 \\ \dots \\ x_n \end{bmatrix}$ with $x_1 = 1 + x_2$ ($n \geq 2$)
- Every vector space with more than one member has at least ____ subspaces.
- Name subspace for $\mathbb{R}^2, \mathbb{R}^3, \mathbb{R}^4$
- Following figures are subspace of \mathbb{R}^2 ?





Example

Let H be the set of all vectors of the form $\begin{bmatrix} a - 3b \\ b - a \\ a \\ b \end{bmatrix}$ where a, b are arbitrary scalars.

That is, let $H = \left\{ \begin{bmatrix} a - 3b \\ b - a \\ a \\ b \end{bmatrix} : a, b \text{ in } R \right\}$. Show that H is a subspace of \mathbb{R}^4 .



Example

- Set of all continuous real-valued functions on \mathbb{R} is a subspace of the vector space of all functions on \mathbb{R} .
- Set of all differentiable real-valued functions on \mathbb{R} is a subspace of the vector space of all functions on \mathbb{R} .
- Set of all functions $D(f(x)) = f'(x)$ is a subspace of the vector space of all functions on \mathbb{R} .

Intersection of subspaces



Theorem

If W_1 and W_2 are subspaces of V , then $W_1 \cap W_2$ is a subspace.

Proof:

$W_1 \cap W_2$ is the largest subspace contained in W_1 and W_2 both.



Theorem

Intersection of any collection of subspaces of a vector space V , is a subspace of V .

Proof:


Union of subspaces



Theorem

The union of two sub-spaces may not a subspace.

Proof:





Theorem

Fact: The union of two sub-spaces is not a subspace unless **one is contained in the other**.

W_1 and W_2 are subspaces of V , then $W_1 \cup W_2$ is subspace of V **if and only if**
 $W_1 \subseteq W_2$ or $W_2 \subseteq W_1$

Proof:

Span and Subspace



Theorem

If v_1, v_2, \dots, v_p are in a vector space V , then $\text{Span} \{v_1, v_2, \dots, v_p\}$ is a subspace of V .

Proof:

Sum of subspaces



- There are two reasons to use the sum of two vector spaces.
 - to build new vector spaces from old ones.
 - to decompose the known vector space into sum of two (smaller) spaces.

- Since we consider linear transformations between vector spaces, these sums lead to representations of these linear maps and corresponding matrices into forms that reflect these sums. In many very important situations, we start with a vector space V and can identify subspaces “internally” from which the whole space V can be built up using the construction of sums.



Definition

Let A and B be non-empty subsets of a vector space V . The **sum of A and B** , denoted $A+B$, is the set of all possible sums of elements from both subsets: $A + B = \{a + b : a \in A, b \in B\}$

Example

- $A = \{(2,3)\}$ $B = \{t(3,1) | t \text{ is scalar}\}$, $A+B$?

- $A = \{t_1(1,2,0) | t_1 \text{ is scalar}\}$ $B = \{t_2(0,1,2) | t_2 \text{ is scalar}\}$, $A+B$?



Theorem

If W_1, \dots, W_m are subspaces of V , then $W_1 + \dots + W_m$ is a subspace of V .

Direct sum of subspaces



Definition

$U + W$ is called a **direct sum**, if any element in $U + W$ can be written uniquely as $u + w$ where $u \in U$ and $w \in W$ (Notation: $U \oplus W$)

Example

Check where sum of following elements is a direct sum?

$$\text{a) } U = \begin{bmatrix} a \\ b \\ 0 \end{bmatrix}, W = \begin{bmatrix} 0 \\ 0 \\ c \end{bmatrix}$$

$$\text{b) } U = \begin{bmatrix} a \\ b \\ 0 \end{bmatrix}, W = \begin{bmatrix} 0 \\ c \\ d \end{bmatrix}$$



Theorem

If U and W are subspaces of V , then the sum is a direct sum $U \oplus W$, if and only if $U \cap W = \{0\}$

Proof:



Example

Let E denote the set of all polynomials of even powers.

$E = \{a_n t^{2n} + a_{n-1} t^{2n-2} + \dots + a_0\}$, and O be the set of all polynomials of odd powers :

$O = \{a_n t^{2n+1} + a_{n-1} t^{2n-1} + \dots + a_0\}$.

The set of all polynomials P is a direct sum of E and O :

$$P = E \oplus O$$

It is easy to see that any polynomial (or function) can be uniquely decomposed into direct sum of its even and odd counterparts:

$$p(t) = \frac{p(t) + p(-t)}{2} + \frac{p(t) - p(-t)}{2}$$



Example

Prove set of all bound functions such as

$$W = \{f(x) \mid \exists M \in \mathbb{R} \text{ such that } |f(x)| \leq M, \forall x \in \mathbb{R}\}$$

is a subspace of $V = \{\text{all functions from } \mathbb{R} \text{ to } \mathbb{R}\}$

Note

Triangle Inequality for Real Numbers

$$|a + b| \leq |a| + |b|$$



- ❑ LINEAR ALGEBRA: Theory, Intuition, Code
- ❑ David Cherney,
- ❑ Online Courses!
- ❑ Chapter 4 of Elementary Linear Algebra with Applications
- ❑ Chapter 3 of Applied Linear Algebra and Matrix Analysis